

CHOLESTERZYME FORMULA

Helps Retain Healthy Cholesterol Levels*



ALCOHOL-FREE CONCENTRATED EXTRACTS OF:

FORMULA		MG EXTRACT PER 2 CAPSULES
Pantethine		200 mg
Artichoke leaf	(<i>Cynara scolymus</i>)	200 mg
Coleus root	(<i>Coleus forskohlii</i>)	160 mg
Guggulu resin	(<i>Commiphora mukul</i>)	150 mg
Arjuna bark	(<i>Terminalia arjuna</i>)	100 mg
Wild Yam rhizome	(<i>Dioscorea villosa</i>)	22 mg
Policosanol		8 mg
Greater Celandine tops & roots	(<i>Chelidonium majus</i>)	8 mg

DOSAGE

2 capsules, 2 times daily

DURATION OF USE

6-12 months

BEST TAKEN

Between meals, with a small amount of warm water

DESCRIPTION OF FORMULA

The Cholesterzyme Formula formula includes herbs and nutrients that function synergistically to help preserve the healthy metabolism of cholesterol, particularly in the liver. Cholesterzyme Formula also protects low-density lipoprotein (LDL) cholesterol from the damaging effects of free radicals. Adopting a healthy lifestyle that includes a daily exercise regimen and a diet high in unprocessed foods (fresh vegetables, fruits and whole grains), will optimize the effects of Cholesterzyme Formula in maintaining optimal cholesterol levels and a vibrant cardiovascular system.*

Guggulu extract is a resinous derivative of a close relative of myrrh that comes from India. Guggulu extract helps to maintain normal cholesterol and triglyceride levels by optimizing cholesterol metabolism in the liver, promoting the excretion of cholesterol through the bile, and protecting LDL cholesterol from the damaging effects of free radical oxidation.*

Policosanol (sometimes called octacosanol) is a wax found in *Saccharum officinarum* (sugarcane). Policosanol helps to maintain a healthy balance between LDL and HDL cholesterol by promoting the proper metabolism of cholesterol in the liver. Accordingly, policosanol plays an important role in maintaining a healthy cardiovascular system.*

Artichoke leaf contains bitter compounds known as caffeoylquinic acids that have multiple beneficial effects on cholesterol metabolism. Artichoke reduces the formation of cholesterol in the liver, promotes the excretion of cholesterol through the bile and protects cholesterol from the damaging effects of free radicals. Collectively, the actions of this leaf serve to promote optimal levels of cholesterol in the body and help to maintain healthy heart function.*

Pantethine is a form of vitamin B5 that helps to maintain normal LDL and HDL cholesterol levels.*

Coleus forskohlii root contains forskolin, a compound that

works directly on the heart muscle and on blood vessels to optimize blood flow throughout the cardiovascular system. Forskolin also helps to maintain healthy fat metabolism.*

Wild yam root contains diosgenin, a constituent that preserves the proper flow of bile, particularly in the presence of estrogen, a hormone that can sometimes reduce the flow. Consequently, this root balances cholesterol levels by promoting the normal excretion of cholesterol through the bile.*

Greater celandine herb is traditionally regarded as an herb that helps to support the healthy flow of bile and for that reason helps to maintain proper cholesterol balance. More research is warranted to validate its use in promoting heart health.*

Arjuna herb is derived from the bark of a tree, which contains several important constituents, including *Terminalia arjuna*, gallic acid and ethyl gallate. These elements help to maintain healthy cholesterol levels and a healthy heart. Likewise, several studies suggest that, when used alone or in conjunction with conventional heart medications, Arjuna helps preserve the proper functioning of the cardiovascular system, and fittingly, the bark is traditionally used in Ayurvedic medicine (from India) to rejuvenate the heart.*

COMPLIMENTARY HERBS/FORMULAS

Cholesterzyme Formula combines well with Hawthorn Supreme.

SAFETY EVALUATION/CONTRAINDICATIONS

This formula should be avoided in pregnancy and lactation. It may occasionally cause an upset stomach, in which case it should be taken with food. It may cause loosening of stools, which is normal and will tend to pass with time. Use with caution if you are allergic to the daisy or chrysanthemum family. If you experience fast or irregular breathing, itching, skin rash or hives, seek medical attention promptly. Also, use with caution and seek the advice of a qualified healthcare professional if you have liver disease, hemophilia or a stomach or intestinal blockage. Do not use for more than three

consecutive months.*

KNOWN DRUG INTERACTIONS

This formula should be used with caution in combination with drugs that inhibit blood clotting and platelet aggregation including but not limited to warfarin (Coumadin®), heparin, clopidogrel (Plavix®), pentoxifylline (Trental®), and aspirin. Additionally, this formula may interact with antihistamine and antihypertension medications. There is also a possibility that this formula might interact with levodopa, a treatment for Parkinson's disease, and theoretically this formula may increase the side effects associated with neostigmine (Prostigmine®), and some eye drops including echothiophate iodide (Phospholine®) and isoflurophate (Fluropryl®). Some of the herbs in this formula have not been studied extensively to determine their interactions with other medications. Before using this formula, talk with your healthcare professional if you take any medications.*

REFERENCES

- Accatino L, Pizarro M, Solis N, Koenig CS. Effects of diosgenin, a plant-derived steroid, on bile secretion and hepatocellular cholestasis induced by estrogens in the rat. *Hepatology* 1998;28(1):129-40.
- Araghiniknam M, Chung S, Nelson-White T, et al. Antioxidant activity of dioscorea and dehydroepiandrosterone (DHEA) in older humans. *Life Sci* 1996;59(11):PL147-57.
- Batista J, Stüsser R, Penichet M, et al. Doppler-ultrasound pilot study of the effects of long-term policosanol therapy on carotid-vertebral atherosclerosis. *Curr Ther Res* 1995;56:906-914.
- Baumann JC, Heintze K, Muth HW. Clinico-experimental studies on the secretion of bile, pancreatic and gastric juice under the influence of phytocholagogous agents of a suspension of *Carduus marianus*, *Chelidonium* and *Curcuma*. *Arzneim Forsch* 1971;21:98-101.
- Benítez M, Romero C, Más R, et al. A comparative study of policosanol versus pravastatin in patients with type II hypercholesterolemia. *Curr Ther Res* 1997;58:859-867.
- Bharani A, Ganguly A, Bhargava KD. Salutory effect of *Terminalia Arjuna* in patients with severe refractory heart failure. *Int J Cardiol* 1995;49(3):191-9.
- Blumenthal M, et al. ed. *The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines*. Trans. S. Klein. Boston, MA: American Botanical Council, 1998.
- Castaño G, Canetti M, Moreira M, et al. Efficacy and tolerability of policosanol in elderly patients with type II hypercholesterolemia: a 12-month study. *Curr Ther Res* 1995;56:819-828.
- Chander R, Khanna AK, Kapoor NK. Lipid lowering activity of guggulsterone from *Commiphora mukul* in hyperlipidaemic rats. *Phytother Res* 1996;10:508-11.
- Dwivedi S, Agarwal MP. Antianginal and cardioprotective effects of *Terminalia arjuna*, an indigenous drug, in coronary artery disease. *J Assoc Physicians India* 1994;42(4):287-9.
- Englisch W, Beckers C, Unkauf M, et al. Efficacy of artichoke dry extract in patients with hyperlipoproteinemia. *Arzneim Forsch* 2000;50:260-5.
- Felter HW, Lloyd JU. *King's American Dispensatory 2 vols.*, 18th ed. Portland OR: Eclectic Medical Publications, 1898, 1983.
- Hoffman BB, Chang H, Reaven GM. Stimulation and inhibition of lipolysis in isolated rat adipocytes: Evidence for age-related changes in responses to forskolin and PGE1. *Horm Metabol Res* 1987;19:358-360.
- Kelly GS. Pantethine: A review of its biochemistry and therapeutic applications. *Alt Med Rev* 1997;2(5):365-77.
- Kraft K. Artichoke leaf extract—recent findings reflecting effects on lipid metabolism, liver and gastrointestinal tracts. *Phytomedicine* 1997;4:370-8.
- Menéndez R, Arruzazabala L, Más R, et al. Cholesterol-lowering effect of policosanol on rabbits with hypercholesterolaemia induced by a wheat starch-casein diet. *Br J Nutr* 1997;77:923-932.
- Ram A, Lauria P, Gupta R, et al. Hypocholesterolaemic effects of *Terminalia arjuna* tree bark. *J Ethnopharmacol* 1997;55(3):165-9.
- Satyavati GV. Guggulipid: A promising hypolipidemic agent from gum guggul (*Commiphora wightii*). *Econ Med Plant Res* 1991;5:47-82.
- Shaila HP, Udupa SL, Udupa AL. Hypolipidemic activity of three indigenous drugs in experimentally induced atherosclerosis. *Int J Cardiol* 1998;67(2):119-214.
- Singh K, Chander R, Kapoor NK. Guggulsterone, a potent hypolipidaemic, prevents oxidation of low density lipoprotein. *Phytother Res* 1997;11:291-4.
- SnowJM. *Coleus forskohlii* Willd (Lamiaceae). *Prot J Botan Med* 1995;Autumn:39-42.
- Thakur CP, Thakur B, Singh S, et al. The Ayurvedic medicines Haritaki, Amala and Bahira reduce cholesterol-induced atherosclerosis in rabbits. *Int J Cardiol* 1988;21(2):167-75.

*THIS STATEMENT HAS NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION.
THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.